

Moly Brook Molybdenum Property, Newfoundland Tenajon Resources Corp.

LOCATION: The Moly Brook Property is located on the south coast of Newfoundland, 1 km north of the community of Grey River. Mean elevation is 250 m.

STORY: Exploration has outlined a 2.5 km long and open trend of molybdenum +/- copper +/- silver mineralization in which three zones, Moly Brook, Wolf and Chimney Pond, are identified as having the potential to host significant, open pitable, molybdenum deposits. The Moly Brook Zone is the most advanced target with forty-four drill holes testing the deposit. The results show the mineralization, alteration and structural characteristics of the zone to be indicative of a relatively shallow-level porphyry-type system associated with an inferred subjacent granitoid stock.

GEOLOGY: The Moly Brook Zone is hosted by deformed Silurian-aged plutonic rocks, ranging in composition from quartz monzonite to granodiorite to locally diorite. The gradational / diffuse contacts between rock types imply derivation from a heterogeneous intermediate-felsic source. At depth, the system is intruded by undeformed granitic dykes containing rare disseminated molybdenum. These dykes cross-cut, and are cut by mineralized veins, indicating both a temporal and genetic relationship to mineralization and are believed to represent the apophyses of a buried stock. Sheeted veining is dominant at the higher elevations passing into a stockwork at depth. Molybdenite typically occurs as fine-grained anhedral disseminations and stringer flakes along vein margins, or is locally disseminated within veins and adjoining microfractures. Chalcopyrite is subordinate to molybdenite and appears more common within Mo-poor, feldspar-rich aplitic veins and later massive quartz-pyrite veins.

DEPOSIT: The Moly Brook Zone has been traced for 750 m along strike, to depths of in excess of 300 m from surface, over widths variable to 500 m. The zone is open along strike and at depth. Drill results include a 378.33 m intercept averaging 0.078% molybdenum. Anomalous copper and silver values overlap and peripheral to the molybdenum deposit. Drill results include an intercept of 127.70 m averaging 0.133% Cu and a 30.49 m intercept averaging 5.39 g/t Ag.

The Wolf Pond Zone hosts a 200 m long x up to 250 m wide of sheeted quartz veining containing highly anomalous molybdenum values with channel samples assaying up to 0.191% molybdenum over 3 m.

In 1965 two short pack sack drill holes were drilled within a 200 m wide x 400 m long molybdenum in soil anomaly near Chimney Pond. Both holes intersected anomalous molybdenum values over their entire length with one hole intersecting a 27.44 m section averaging 0.057% molybdenum, the last 7.62 m of which averaged 0.118% molybdenum.

DISPLAY: Core box containing samples from the Moly Brook Zone, cross sections and maps to illustrate the property will be displayed.