

Rouyn Gold Exploration Property

Yorbeau Resources, Inc.

LOCATION: The Property is located 5 km south of the town of Rouyn-Noranda in northwestern Quebec, Canada. Mean elevation is 250 m.

STORY: The Rouyn Property lies within the Abitibi Greenstone Belt of the Superior Province and occurs in the Rouyn-Val d'Or segment that hosts numerous gold deposits along the Cadillac-Larder Lake Break (the "CLLB"), a major fault zone that is one of the truly fundamental structures in the Late Archean Abitibi Greenstone Belt. Gold values are widespread on the Rouyn Property and they are mainly found associated with the CLLB in rock lithologies belonging to the Piché Group.

GEOLOGY: The Property straddles the CLLB along an E-W direction for 12 km. The oldest rocks in the immediate area are schists and migmatites belonging to the Pontiac Group. They are overlain by conglomerates, sandstones and siltstones of the Temiscaming Group. This group is capped by the CLLB rocks comprising carbonate rocks, talc-chlorite and chlorite, and minor sandstone interbeds. The Temiscaming Group is in contact to the north with the Blake River Group. The general strike on the Augmitto and Cinderella blocks is E-W. Dips are north 70-80°.

DEPOSITS: Gold values are widespread on the Rouyn Property. The major gold occurrences are associated with the CLLB. Both the Augmitto and Astoria deposits occur on the CLLB. The Lac Gamble and Cinderella Zones contain thick zones of silicification, the strongest type of alteration associated with the gold mineralization, in the Piché Group. The relative sequence of alteration from the least altered komatiitic rocks is: 1) talc-chlorite (\pm calcite) schist followed by; 2) "green carbonate" rocks (carbonatized komatiites with fuchsite), which is the lower level of alteration related to gold deposition (generally low gold grades, < 3 g/t), which is superimposed by the most intense alteration; 3) silicification and quartz veining, frequently accompanied by brecciation. Higher levels of gold occur in these silicified zones with minor arsenopyrite, carbonates of calcium, iron and magnesium and brown tourmaline.

DISPLAY: Four core boxes displaying drill core from the Augmitto deposit, Lac Gamble and Cinderella targets are available for review. Examples of gold contents include: Lac Gamble hole 461 (10.0 m core length @ 13.77 g/t Au); Lac Gamble hole 457 (4.5 m @ 9.84 g/t); and Cinderella hole 467 (4.8 m @ 27.3 g/t). Cross sections and maps will illustrate the mineralization.